

---

**POSTPARTUM INTERVAL OF SOME LOCAL BREEDS OF COWS  
IN SOUTH DARFUR STATE - SUDAN**

**BY**

*MOHAMMED YOUSIF ABDALLA ALFAKI*

*B. V. Sc. ZAGAZIG UNIVERSITY*

*1982*

***SUPERVISOR: DR. ADIL SALIM ELSHEIKH***

***Ph. D. HOKKAIDO***

**A THESIS SUBMITTED IN PARTIAL FULFILLMENT  
OF THE REQUIREMENTS OF UNIVERSITY OF  
KHARTOUM FOR THE MASTER DEGREE  
OF VETERINARY MEDICINE**

**Department of Reproduction and Obstetrics  
Faculty of Veterinary Medicine  
University of Khartoum**

**February 2006**

## **Dedication**

*To my late father, my merciful,*

*Mother, brothers, sisters.*

*To my beloved wife Amal Mahmood*

*To the lovely eyes of my daughters*

*Soso, Yasmin, Zoba ,Tota and Razan.and also*

*To my relatives and friends.*

## **Acknowledgements**

I thank God who gave me the spirits, strength and patience to achieve my goal.

I want to express my sincere appreciation and deepest gratitude to my supervisor Dr. Adil Salim Elsheikh for his contact, encouragement, valuable advises, generous support and appreciable guidance at all stages of writing to complete these work. I am indebted for him for supervision, unlimited help and patience during this study, which made it possible for me to complete this work.

Especial thanks are extended to all staff members of the Department of Reproduction and Obstetrics in the Faculty of Veterinary Medicine. U of K.

I wish to express my warm thanks and special gratitude to my best friends Dr. Mohammed Salih Mousa, Dr. Faisal Omer, Dr. Basheer Abdalla and Dr. Shiekh Eldin Ahmed Mahmoud for their continuous encouragement and logistic support.

My utmost appreciation and gratitude to my uncle Ibrahim Abdalla Alfaki and to my Sister Haga Thorya Mahmoud for their encouragement.

I express my thanks to stockowners Mousa Alsharief, Abdalla Masar, Osman Magid, Hassan Ahmed Ismail and Omer Atalla for their facilities to collect the data for this study.

Also I wish to express my deepest thanks to Mr. Ibrahim Aldalil the minister of Regional Ministry of finance and Dr. Bahar Aldin Salih Ibrahim the minister of the regional ministry of Agriculture and Animal Resources for their financial support of this study.

My special thanks are also due to Mr. Ahmed Rabih the head of Human Resources Department in South Darfur State.

Finally I wish to thanks Mrs. Yousra Elmagzoub for her patience during typing this thesis.

## TABLE OF CONTENTS

<i>No</i>	<i>Subject</i>	<i>Page</i>
1	Dedication	I
2	Acknowledgment	II
3	Table of contents	III
3	List of figures	VI
4	List of Tables	VII
5	Abbreviations	VIII
6	List of Appendices	X
7	Abstract	XI
8	Arabic Abstract	XIII
9	Introduction	1-5
<b><i>Chapter -1- Literature Review</i></b>		
1.1	Oestrous cycle	6
1.2	Signs of oestrous	6
1.3	Silent heat or suboestrus	7
1.4	Postpartum interval	8
1.4.1	Definition	8
1.4.1	Postpartum changes	9
1.4.2	Uterine involution	9
1.4.2.1	Regression of endometrium	9
1.4.2.2	Lochia (Fluid discharge)	9

1.4.2.3	Elimination of bacterial contamination	10
1.4.2.4	Resumption of the ovarian activities	10
1.4.3	Factors affecting PPI	11
1.4.3.1	Nutrition	11
1.4.3.2	Body condition score (BCS)	11
1.4.3.3	Suckling	12
1.4.3.4	Age	13
1.4.3.5	Season	13
1.4.3.6	Hormonal activity during postpartum	14
1.4.3.7	Milking	17
1.4.3.8	Heat stress	17
1.4.3.9	Uterine infections	18
1.4.3.10	Management	19
1.4.3.11	Breed	19
1.4.3.12	Biological uses of GnRH	19
1.4.4	Calve interval	21

## ***Chapter -2- Materials and Methods***

2.1	Study area	23
2.2	Climate	23
2.3	Temperature measurement	23
2.4	Animals	24
2.4.1	Management and feeding	24
2.4.2	Housing	25
2.4.3	Milking	25
2.4.4	Pregnancy diagnosis	25

2.4.5	Health control	26
2.5	Experiment design	27
2.5.1	Experiment (1)	27
2.5.2	Experiment (2)	27
2.5.3	Experiment (3)	28
2.6	Statistical analysis	28

### ***Chapter -3- Results***

3.1	Experiment 1	29
3.1.1	The effect of the breed on the length of the PPI	29
3.1.2	The relation between the BCS and the PPI	29
3.1.3	The relation between the parity and the PPI	29
3.2	Experiment (2)	33
3.3	Experiment (3)	37

### ***Chapter -4-***

4.1	Discussion	39
4.2	Conclusion	44
4.3	Recommendation	45
	References	46
	Appendices	65

## LIST OF FIGURES

<i>Figure</i>	<i>Title</i>	<i>Page</i>
1-	Postpartum interval in three different dairy breeds-bred in South Darfur State.	30
2-	The relation between the BCS and the length of the PPI of the local breeds Kenana, Fellata and cross-bred dairy cows.	31
3-	The relation between the parity and the length of the PPI of the local breeds Kenana, Fellata and cross-bred dairy cows.	32
4-	The effect of GnRH treatment on day 21 postpartum on the length of PPI of Kenana cows	34
5-	The effect of GnRH treatment on day 21 postpartum on the length of the PPI of Fellata cows	35
6-	The effect of GnRH treatment on day 21 postpartum on the length of the PPI of cross-bred cows	36
7-	The effect of the season on the length of the PPI of the three breeds, Kenana, Fellata and cross-bred dairy cows	38



## **List of Tables**

<b>Table</b>	<b>Title</b>	<b>Page</b>
1.	Population of livestock in South Darfur State.	2
2.	The average rain fall in Nyala Metrological Station.	

24

## **ABBREVIATIONS**

APU	Agriculture planning unit
BCS	Body condition score
BQ	Black quarter
CBPP	Contagious bovine pleuro pneumonia
CBS	Central Bureau of Statistic
CI	Calving interval
CL	Corpus luteum
COA	Cyclic ovarian activity
CRH	Corticotrophin releasing hormone
DF	Dominant follicle
FAO	Food Agriculture Organization
FSH	Follicle stimulating hormone
FSH-RH	Follicle stimulating hormone releasing hormone

GH	Growth hormone
GnRH	Gonadotropic-releasing hormone
HS	Haemorrhagic septisemia
LH	Luteinizing hormone
LH-RH	Luteinizing hormone releasing hormone
MAAR	Ministry of Agriculture and Animal Resources
PGF <sub>2</sub> $\alpha$	Prostaglandin F <sub>2</sub> $\alpha$
PGS	Prostaglandins
PIF	Prolactin inhibitory factor
PPI	Postpartum interval

## **List of Appendices**

<b>Appendix</b>	<b>Title</b>	<b>Page</b>
1	Annual rain fall	65
2	Population of livestock in South Darfur State	66
3	Population data sheet	67
4	Average temperature	68
5	Different Parameters of PPI of local dairy cows native to South Darfur State	69
6	PPI of local breeds of dairy cows native to South Darfur State treated with GnRH	71
7	The length of PPI of dairy cows native to South Darfur State during the three different seasons	72
8	The effect of season of birth on the PPI of Kenana, Fellata and cross-bred dairy cows	73
9	PPI of control Kenana dairy cows	73
10	PPI of control Fellata dairy cows	74
11	PPI of control cross-bred dairy cows	75
12	Treated Kenana dairy cows	76
13	Treated Fellata dairy cows	76



## ABSTRACT

In the present study, the length of postpartum interval (PPI), the effect of season and gonadotropin (GnRH) injection on the length of PPI were studied in some local Sudanese dairy cows native to South Darfur State namely Fellata, Kenana and cross-bred. Furthermore the influences of body condition score (BCS) and parity on PPI were also investigated.

In experiment 1 a total of 59 dairy cows that gave birth were employed to determine the length of PPI. The parity range of these cows was 1 to 5. Their BCS was between 2.5 to 4. The cows were grouped according to their breeds into three groups. Group I was Kenana cows ( $n = 21$ ), group II was Fellata cows ( $n = 17$ ) and group III was cross-bred cows ( $n = 21$ ). The PPI was recorded as the time elapse between parturition and the appearance of first oestrus postpartum. The pure local breeds namely Kenana and Fellata cows had a significantly ( $P < 0.001$ ) longer PPI compared to cross-bred dairy cows. The PPI of Kenana, Fellata and cross-bred cows were  $286.9 \pm 35.3$  days,  $246.3 \pm 26.6$  days and  $122.00 \pm 14.9$  days respectively. No differences ( $P > 0.05$ ) in the PPI were found between Kenana and Fellata cows. Also a slight negative relation ( $R = 0.2$ ) between BCS and the PPI was observed. Parity has no influence ( $R = 0.005$ ) on PPI, however when parity increases the PPI decreases.

Experiment II was designed to determine the effect of GnRH on the PPI of the above mentioned breeds of cows. A total of 66 dairy cows (Kenana  $n = 22$ , Fellata  $n = 22$ , cross-bred  $n = 22$ ) having the same BCS and parity as those in experiment I were used in this experiment. The cows were grouped into two groups. Each group was 33 cows (Kenana  $n = 11$ , Fellata  $n = 11$ , cross-bred  $n = 11$  cows). Group I was intramuscularly injected with

250 µg of GnRH on day 21 postpartum. Group II was left untreated to serve as control. GnRH treatment significantly ( $P < 0.001$ ) reduced the PPI in all breeds. The PPI of the treated Kenana cows, Fellata cows and cross-bred cows was  $79.5 \pm 8.4$ ;  $65.9 \pm 3.2$  days and  $74.2 \pm 2.8$  days respectively. No differences ( $P > 0.05$ ) in PPI among the treated breeds was observed.

Experiment III was designed to study the influences of season of birth on the PPI of the above mentioned cows. The PPI was compared among three groups of cows that gave birth in three different seasons. Group I was 18 cows that gave birth in Summer (Kenana  $n = 7$ , Fellata,  $n = 4$  and cross-bred  $n = 7$ ). Group II was 19 cows that gave birth in Autumn (Kenana  $n = 9$ , Fellata  $n = 6$  and cross-bred  $n = 4$ ) and group III was 31 cows gave birth in Winter (Kenana  $n = 13$ , Fellata  $n = 11$  and cross-bred  $n = 7$ ). The BCS and parity of these cows were similar to those of the above experiments. The results showed that the PPI of cross-bred cows was significantly ( $P < 0.001$ ) shorter in Summer. No difference ( $P > 0.05$ ) in PPI of cross-bred cows that gave birth in Winter and Autumn was observed. Neither the PPI of Kenana nor that of Fellata cows were influenced by season ( $P > 0.05$ ). It is concluded that the mean PPI of Kenana, Fellata and cross-bred cows in South Darfur State were  $286.9 \pm 35.3$  days,  $246.3 \pm 26.6$  days,  $122.00 \pm 14.9$  days respectively. GnRH injection on day 21 postpartum reduces the PPI of the three breeds of cows. Additionally, the PPI of the cross-bred cows is only influenced by Summer season. However season has no effect on the PPI of pure local cows.

(season)	(PPI)	(GnRH)	
	(BCS)		.
		.(PPI)	(Parity)
PPI		59	
		.PPI	
4- 2.5			5-1
( 21 = )			.
.( 21= )			( 17= )
.			
		(	)
122.00 ± 14.9,			.(p<0.001)
			246.3 ± 26.6 , 286.9 ± 35.3
			(P>0.05)
			(R=0.2)
			(R= 0.005)
			.
PPI	GnRH		
22=	22= )	66	.
parity	BCS		.( 22=
11= )	33		.



GnRH	250 μg	11=	11=
(p<0.001)		21	
79.5 ± 8.4			
(P>0.05)		74.2 ± 2.8	65.9 ± 3.2
	(season)		
4=	7=	18	
6=	9=	19	7=
13=		31	4=
			7=
			11=
(P>0.05)			(p<0.001)
		(P>0.05)	
122.0 ±14.9	246.3 ± 26.6	286.9 ± 35.3	
	21	GnRH	

## **INTRODUCTION**

South Darfur State of Western Sudan comprises 135000 square kilometers of semi arid and savannah zones, and lies within the area defined by the coordinate, Latitude 13° - 9:30 north and Longitude 27° - 24:30 east. It has common boarders with North Darfur, West Kordofan, West Darfur and also North Baher Elgazal State. It also has a shared boarder with Chad and Central Africa Republics.

The human population of South Darfur State is about 3.171 millions (CBS, 2004), who are mainly traditional farmers and livestock owners (80% of the total population) live stock producers in the area are nomad, transhumant and sedentary. Nomads and transhumant annually move with their animals from South to North of the state at the beginning of the rains avoiding biting insects and mud in the South. The live stock provides food and capital for human welfare and economic development. South Darfur State is one of the richest states in the country; it is well famous with the natural resources according to its geographical location between the poor Savanna to the North, and the rich Savanna to the South, as well as the Mediteranian climate of Gabal Mara area.

The annual rainfall lies between 300 - 900 mm a year increasing from North to South (Meterological office Nyala station, appendix .1.). The livestock population is estimated as 10 million head according to the Regional Ministry of Agriculture and Animal Resources figures (1993), table .1. and appendix .2.

Table .1. Population of livestock in South Darfur State

SPECIES	No	Percentage/ total population of the Sudan
Cattle	3.476.516	10% (34.276.616)
Sheep	3.548.022	8.37% (44.350.275)
Goats	2.519.445	6.9% (35.992.071)
Camels	69000	2.3% (3.019.348)
Equines	300.000	3% (10.000.000)

South Darfur State is the richest state in the Sudan according to the population of animals. The cattle in this state have special properties which became very famous locally and nationally, it is known as Nyala Cattle, or Nalaweyan cattle referred to the name of the capital of the state (Nyala).

The cattle in this state originate from the zebu cattle which is known in the Sudan as Buggara cattle which has the following properties: -

1. It is a dual purpose cattle for meat and milk production.
2. It is highly resistant to the diseases specially tick born diseases.
3. It has excellent resistance against the external factors and can move for long distance for food and water.
4. It has the ability to stay in very bad conditions and under poor feeding.
5. It has the ability to increase rapidly in weight under favorable conditions.

There are different types of breeds of cattle in the state including:-

1. Local breeds which are known as Nalaweyan cattle.
2. Kenana breed.
3. Cross-bred.

The Local breeds in the state including:-

A. Rizagat cattle: which are red in colour, small in size, with less milk production and can move for long distances.

B. Fellata cattle: It is mainly white in colour, large in size and comparatively with more milk production. This breed is owned by Fellata tribe, Banihalpa, Taisha, Targam, Birgid, and Dago, tribes.

Kenana breed is introduced firstly to the state by AL Gazala Gawazat Research Center to improve the local breeds around the project area 1964, (Sideg Ibrahim Yahia) (personal communication).

Cross-breeds were introduced recently (1991) to the state specifically in Nyala City by Mr. Kamal Adeep and in Mara Mountain, by Mr. Dirag. (personal contact) The cross breeds are mainly kept for milk production. They were a result of crossing between imported Friesian bulls with the local Kenana breed in the state. Recently there is a trend to improve the local breeds for milk and meat production.

In South Darfur State the dairy cattle constitute a very small proportion and the rest are dual-purpose. With such small population of dairy cattle, the production of milk and its products for consumption does not meet the demand. To meet this demand, regular calving at an optimal Interval of about 365 days is desirable as it results in more calves being

born, increase the life time, milk production per cow and maximize income (Galina and Arthur, 1989).

Although the indigenous zebu cattle are more adapted to the local tropical environment, their capacity for milk production is usually low (Vaccaro, 1974). Selection for high milk production within the indigenous cattle requires long term genetic improvement programme.

Ageeb, and Hillers (1991) reported that, the indigenous breeds in developing countries generally produce relatively little milk, and there are additional important constraints affecting the production like, quality/quantity of feed, direct and indirect climatic factors and husbandry practices.

It is generally believed that a calving interval of 12 months is an ideal goal for economic dairy cattle production. This goal can not be achieved unless that postpartum interval (PPI) to first oestrus in dairy cows is reduced. The zebu cattle namely *Bos indicus* possess a long PPI (Osman and Eladin, 1971). The local dairy cows native to South Darfur State belongs to this group of cattle. However sparse information about the length PPI of dairy cows in the Sudan is available (Elzubeir, 2003; Elhag, 2003; Elsheikh and Ahmed 2005). This information was for cross-bred dairy cows only. Unfortunately the PPI of the pure local dairy cows is not authenticated in the Sudan. Therefore the objectives of the current study were:-

A. The general objectives were:-

1. To determine the length of the PPI of the local dairy breeds of cattle in South Darfur State, including the Fellata breed, Kenana breed and cross-bred.
2. To determine the effect of parity and BCS on the length of the PPI.

3. To determine the effect of GnRH injection on day 21 postpartum on the length of the PPI.
4. To study the effect of the season of parturition on the length of the PPI in the above mentioned breeds.

B. The main objectives were:-

1. To reduce the length of the PPI of the local dairy breeds of cows in South Darfur State of Western Sudan, aiming to make the cows more productive giving a calf every year, to be more economic and to give more milk to face the demand of the fresh milk specially in the dry season where the nomads and semi nomads move with their cattle to the South and Western parts of the state.
2. To implement the regimen described by Elzubeir and Elsheikh (2004) for management of postpartum period by injecting GnRH on day 21 postpartum.

# **1. Literature Review**

## **1.1 Oestrous cycle**

Oestrous cycle is divided traditionally into five phases.

1. pro-oestrous phase: it is the phase preceding oestrus. It is characterized by marked increase in activity of the reproductive system. There is follicular growth and regression of the corpus luteum of the previous cycle.
2. The oestrous phase: it is the period of acceptance of the male. The female seeks out male and stand for it to mate her. The ovulation occurs during this phase in all domestic animal species except the cow, where it occurs about 12 hours after the end of oestrus. The main hormone produced during oestrous phase is oestrogen.
3. Metoestrous phase: this is characterized by luteinization of the granulosa cells of the ovulated follicle to form the corpus luteum. During this phase there is reduction in the amount of secretions.
4. Dioestrous phase: during this phase the corpus luteum is fully functional and secreting large amount of progesterone.

5. Anoestrous phase: during this phase the genital system remains quiescent, and this phase is only recognized in seasonal breeders.

## **1.2 Signs of oestrus**

Great variations in the intensity of heat signs exist among cows. The manifestations of heat signs tend to be more marked in heifers than in cows. The cow is said to be in oestrus when it stand to be mounted by other cows or bulls (Arthur, *et al.* 1998).

The cow on heat is restless and it shows evidence of bellowing (Arthur, *et al.* 1998). There is a reduction in milk production as results of a reduction in the time spend for eating, rumination and resting (Hafez, 1993). The same author reported that: the oestrous cow licks and sniffs the perineum of other cows, jumps on cows and mounts the bull before standing to be mounted. There is a transparent clear genital discharge from the vulva. The cow shows evidence of pelvic thrusting after full response to the bull and stand with arched back and raised tail. This posture indicates that mating has occurred. In order to improve production in cows it is necessary to achieve a 12 month calving interval. To reach this goal, cows must cycle and become pregnant by an average of 85 days postpartum (Galina and Arthur, 1989; Dimmick, *et al.*, 1991). It was reported that long postpartum anoestrus over 120 days is a very common reproductive problem in primiparous cows under tropical environment (Elhag, 2003; Elshiekh and Ahmed 2005). Although cattle are polyoestrus through out the year, the interaction of the calf with the mother cow was reported to delay the onset of the oestrous cycle (Vicker, *et al.*, 1989). It was also reported by Williams (1990) that temporary weaning of calves reduces the interval to the first oestrus.



The duration of oestrus can be influenced by the breed of the animal, season of the year, presence of bull, level of nutrition, milk yield and number of cows that are in oestrus at the same time (Arthur, *et al.* 1998).

### **1.3 Silent heat or Suboestrus:-**

Some cows and heifers fail to show the signs of oestrus, at the same time they have normal cyclical activity, this condition is known as silent heat or suboestrus. This may be due to failure of the herdsmen to observe the signs of the oestrous cow. The trained staff plays a major role in dairy farms to detect the oestrous cow (Esslemont and Kossaibati, 2000). The first and the second ovulation postpartum may not be accompanied with behavioral signs of oestrus (Arthur *et al.* 1998). The silent heats are very common in Winter than in Summer months. The postpartum anoestrus in cows is reported to be due to failure of ovulation (Savio, *et al.* 1990).

### **1.4 Postpartum Interval (PPI)**

#### **1.4.1 Definition:**

Generally it is known as the period from the day of parturition till the appearance of the signs of the first oestrus postpartum. It is also defined by Malvern (1984) as the period required for recovery of the brain, pituitary, ovaries and genital tract functions after pregnancy and parturition. It was found that 90% of dairy cows start oestrous cycle by 60 days after calving which is the optimal period for the reproductive system to restart its activities for the oestrous cycle, conception, pregnancy, and to give a calf every year (Youngquist, 1988). The time elapsed from parturition until the first postpartum oestrus is accompanied by ovulation and it could be as short as 15 days or longer than 100 days (Malvern, 1984; MacMillan and

Thatcher, 1991). A long PPI over 120 days is a common reproductive problem under tropical condition, (Elhag, 2003; Elsheikh and Ahmed, 2005). The major problem that limits the improvement of reproductive efficiency of dairy cows reared under tropical conditions was the long PPI (Short, *et al.* 1990; Williams, 1990). The Sudanese cross-bred dairy cows have along PPI a round  $162.90 \pm 2.70$  days (Elhag, 2003). The prolonged PPI is attributed to the absence of LH pulses, which lead to atresia of the developing follicles postpartum (Yavas and Walton, 2000). The same authors reported that injection of GnRH early postpartum can initiates the postpartum ovarian cyclicity.

Herdmen and Shepards in South Darfur State observed that the length of the PPI of the local breeds of cows kept under range conditions are above the optimum rang. Also dairy cows in South Darfur Cities were observed to have a long PPI.

### **1.4.2 Postpartum changes**

#### **1.4.2.1 Uterine involution:-**

Uterine involution is defined as the reduction in the size of the genital tract to its normal non-pregnant size and function after parturition. The diameter of the previously gravid horn is halved by five days and it's length is halved by 15 days (Arthur *et al.* 1998).

In this phase there is uterine discharge and the time taken for complete uterine involution ranged from 26-52 days in dairy cattle (Michiel, *et al.* 1999). The uterus return to it is pregravid size within 3-7 weeks depending on the myometral contraction, elimination of bacterial infection and regeneration of the endometrium. (Arthur, *et al.* 1998)

The cervix constricts rapidly postpartum, within 10-12 hours after normal calving and by 96 hours can admit only two fingers (Arthur, *et al.* 1998).

#### **1.4.2.2. Regression of endometrium:-**

Arthur, *et al.* (1998) reported that there is early necrotic changes in the septal mass of the caruncle within the 48 hours postpartum. The same authors mentioned that, complete regression of endometrium occurs soon after parturition in undamaged areas.

#### **1.4.2.3 Lochia (Fluid Discharge):-**

It is the fluid discharge from the uterus or the genitalia postpartum, it is composed of blood, serum mucous, desquamated epithelium, parts of the fetal membranes, parts of the fetal fluids and leucocytes. It is mucoid in consistency and it is yellowish brown or reddish brown in colour (Hafez, 1993). The discharge of lochia starts on the 2<sup>nd</sup> day postpartum and it reaches the maximum on 6 to 8 day postpartum and then decreases and disappears 14 to 18 days postpartum (Hafez, 1993). The whole puerium in cattle takes 42 days, during this period the genitalia retain to it is physiological condition before pregnancy (Arthur, *et al.* 1998). The high level of PGF<sub>2</sub> $\alpha$  after parturition increases the uterine contraction, decreases the uterine size, hasten expulsion of the lochia and expulsion of fetal membranes (Hafez, 1993).

#### **1.4.2.4 Elimination of bacterial contamination:-**

During the last stage of pregnancy there is small widening of the cervical canal especially when the animal lay down, the pathogenic and non

pathogenic bacteria can enter the uterus and rapidly multiply in the favorable media of the uterus. The elimination of the bacteria occurs through phagocytosis and by expulsion of uterine fluids and uterine membranes debris (Hafez, 1993).

#### **1.4.2.5 Resumption of the ovarian activities:-**

The high level of progesterone secreted by the CL and the placenta during pregnancy exerts continuous feed back effect on the pituitary secretions. After parturition the cow enters a period of postpartum anoestrus, although the progesterone level drops down. The duration of the postpartum anoestrus ranged from 30-67 days for dairy cows, the mean interval from calving to first oestrus postpartum ranged from 33-85 days (Arthur, *et al.* 1998).

#### **1.4.3 Factors affecting PPI**

The length of the PPI depends mainly on the level of nutrition, body condition score (BCS), suckling, age, season of birth hormonal influences, milking, heat stress and uterine pathology (Short *et al.* 1990; Stagger *et al.* 1998). Early establishment of cyclic activity after calving is essential for maintenance of an acceptable calving to conception interval (Benmard and Stevenson, 1986). Also it was found that dairy cows ovaries rebound within 30 days (lamming, *et al.* 1981). Some dairy cows may ovulate without showing the oestrous behaviors which lead to long PPI (Hafez, 1993).

#### **1.4.3.1 Nutrition:-**

Nutrition is also among the primary factors that influence the PPI (Mukasa, *et al.* 1991). The metabolic demand for high milk yielding cows should be considered as it has a possible effect on postpartum ovarian activity (Coppock, *et al.* 1974; Butler, *et al.* 1981; Sejresan and Sorensen, 1982; Elkeraby and Abu-Ela 1982). Strategic management of nutrition is also essential for improvement of the reproductive performance of cows (Short *et al.* 1990; Williams, 1995). The length of the postpartum and the reproductive performance of the dairy cows can be affected directly by deficiencies of energy, (Youngquist, 1988). The occurrence of the first postpartum oestrus is influenced by the level of nutrition (Kovacevic, *et al.* 1988). The cows under high level of nutrition during the postpartum period have high incidence of oestrus and ovulation as a result of high level of FSH that increases during follicular phase (Rhind and McNeilly, 1986).

#### **1.4.3.2 The Body condition score (BCS)**

Paul (1995) reported that the BCS involves tactile and visual appraisal of the amount of tissues that cover the vertebrae at the lumber region and around the tail head. Veerkamp, (1998) suggested that successive measures of BCS may be useful indicator of energy balance as the short fall in energy obtained from food is believed to come from mobilization of body tissue reserves.

The BCS is used as a subjective method to determine the body reserves of dairy cattle (Paul, *et al.* 1995) the method is based on a visual

and tactical appraisal of body fat reserves in the back and pelvic regions. The BCS is usually scored on a scale of 1 to 5. (Wildman, *et al.* 1982).

BCS changes over early lactation, could affect the resumption of oestrous cycles and reproductive success. Loss of BCS has been found to be associated with increased levels of milk production at the phenotypic level ( Pryce, *et al.* 2001).

The cumulative percent of cows exhibiting oestrus by day 20, 40 and 60 postpartum was higher when the cows calved at a BCS of scale grade 4 (Richards, *et al.* 1982). When the level of nutrition was high the cumulative percentage of cows exhibiting oestrus during postpartum period was high. If the level of nutrition is increased during postpartum period the level of FSH increases during follicular phase and results in high ovulation rate (Sejresan and Sorensen, 1982).

#### **1.4.3.3 Suckling:-**

Suckling was reported as one of the primary factor that determine the length of PPI in cattle (Mukasa, *et al.* 1991). Suckling induces suppression of LH through suppression of LHRH secretion (Lamming, *et al.* 1982; Acosta, *et al.* 1983; Webb, *et al.* 1996).

Stagger, *et al.* (1998) had shown that restriction of suckling to once daily or complete isolation of calf from suckling beef dams induced an earlier ovulation as compared with continuously suckling dams. Restricted suckling was reported to hasten the return to oestrus. The postpartum pituitary responsiveness and ovarian activity are reported to return within 4-5 weeks after calving in beef cows (Mc Natty, 1988). Strategic management of suckling should improve the reproductive performance of cows without

decreasing the growth rate of calves (Randel, 1981; Short, *et al.* 1990; Williams, 1990; William and Griffith).

#### **1.4.3.4 Age:-**

The PPI of old cows tend to be longer than that of heifers and middle age cows (Marion and Gier, 1998). The same authors reported that primiparous dairy cows have longer PPI than the multiparous. The cows reaching their seventh lactation are known to had a poor fertility (Esslemont and Kossaibati, 2000). However, no effect of age on PPI was reported by (Zainelabdin, 2001).

#### **1.4.3.5 Season:-**

The season has strong effect on the interval between parturition and first oestrus in dairy cows, (Peters and Riley, 1982). It is also found that the interval between calving and first oestrus was longer in Winter and early Spring (Montgomery, 1985). The animals calved in dry Summer season has long calving interval compared to those calved at rainy season.

The breeding season is influenced by environmental and physiological factors that regulating the breeding season. The rain fall and temperatures are considered as environmental factors affecting the season of breeding. Also photoperiodic variations are considered among the major environmental factors that affect the reproductive efficiency. It was found that frequency of oestrus and ovulation, as well as conception rate in Kenyan zebu cattle is higher during the Summer than during the Winter (Rhodes, *et al.* 1982).

#### **1.4.3.6 Hormonal activity during postpartum:-**

- **FSH:-**

The level of FSH increases in concentration from preparturition to reach high levels postpartum (Lamming, *et al.* 1982). This was explained by (Hafez, 1993) by the variable follicular growth and atresia that occur in the preovulatory PPI. The follicular growth with development of mature follicles is prerequisite for LH response and ovulation (Peters and Lamming 1984). Plasma concentration of FSH is also reported to increase during the first 3-5 days postpartum (Lamming, *et al.* 1981; Lamming, *et al.* 1982).

- **LH:-**

Pituitary LH content and responsiveness to GnRH increases rapidly from their lowest at parturition and increases gradually when follicles are recruited postpartum (Malvern, 1984; Schallenberger, *et al.* 1984). The recovery of pituitary LH reserve and its responsiveness to GnRH seems to be nearly completed by day 10 postpartum in dairy cows (Moss, *et al.* 1985). Administration of GnRH stimulates the pituitary gland to induce LH pulses and the LH surge which is responsible for ovulation of the mature follicle (Hafez, 1993).

- **Oestrogens:-**

The level of oestradiol increases from day 26 before parturition and then decreases from day 9 postpartum (Smith, *et al.* 1987). Also oestradiol is necessary for sensitization of pituitary to release both preovulatory FSH and LH. Arthur, *et al.* (1998) reported that oestrogen and progesterone decline from high values during late pregnancy to a very low concentration over the first 4-5 days postpartum. Ginther, *et al.* (1996) reported that the high level



of oestrogen and progesterone concentration prior to calving suppresses the follicular waves before calving.

- **Progesterone**

The plasma concentration of progesterone hormone is high during last stage of pregnancy, and then begins to drop in concentration before parturition and remains very low over the first 4-5 days postpartum.

The sudden drop of progesterone following parturition lead to removal of the negative feed back on the gonadotrophins and thus exerts appositive feed back upon the hypothalamus to release GnRH which stimulates the anterior pituitary to release FSH and LH (Savio, *et al.* 1990). It was found that, the progesterone hormone enhances the luteal phase of the next cycle and controlling oestrous cycle (William, *et al.* 1982; Bashir, 1990).

- **Oxytocin:-**

The level of oxytocin reach its maximum level (6.4pg/ml) at the expulsive stage of parturition and falls to the minimum level (0.5pg/ml) at time of oestrus (Hafez, 1993). The oxytocin level during late pregnancy and early stages of parturition remains low and increases to reach the peak values when the fetal head emerges from the vulva. So it plays only minor role in initiation of uterine contractions. The release of oxytocin hormone is stimulated by sensory receptors in the cervix and anterior vagina.

- **Prostaglandins (PG,s):-**

The level of prostaglandins metabolite were high at parturition and they were elevated on the 7 to 23 days postpartum. PG play an important role in uterine involution postpartum (Lindel, *et al.* 1982). PG specially PGF<sub>2</sub>α are

essential for ovulation and activation of the collagenases that cause weakening of the follicular wall and thus enhances ovulation.

Injection of exogenous PGF<sub>2</sub>α during the 1<sup>st</sup>, the 2<sup>nd</sup>, or the 4<sup>th</sup> week postpartum enhanced the uterine involution of dairy cows (Kindahl, *et al.* 1982). Also it was found that administration of PGF<sub>2</sub>α during the 3<sup>rd</sup>, the 5<sup>th</sup> or the 8<sup>th</sup> week postpartum stimulates early cyclicity in dairy cows (Zemjanis, 1980). Sudanese cross bred cows (Kenana × Friesian) which were given a single or a double injection of PGF<sub>2</sub>α during the 1<sup>st</sup>, 2<sup>nd</sup> or the 3<sup>rd</sup> week postpartum, expressed a shorter postpartum period (Elzubeir and Elsheikh 2004). Some authors reported that a single or double injection of PGF<sub>2</sub>α during the postpartum period at 10-14 days interval decreases the days open (Youngquist, 1988; Schofield, *et al.* 1999).

- **Prolactin :-**

Prolactin increases 2-4 days before parturition and continues to increase after parturition in both milking and suckling cows (Hafez, 1993). It had been reported that suckling and milking had no effect on prolactin levels (Smith, *et al.* 1987). It was also found that the release of prolactin is under the control of prolactin inhibitory factor (PIF) which is released from the hypothalamus. Prolactin is also an important luteotrophic agent.

- **Insulin:-**

Insulin has a gonadotrophic effect on follicular development. This gonadotrophic effect is exerted through its action on the hypothalamic neurons (Lucy, *et al.* 1992; Gong, *et al.* 2002; Gong, *et al.* 1996). In their

report they mentioned that insulin has specific receptors located in the hypothalamus. Insulin binds to these receptors and enhancing glucose metabolism inside the neurons. Thus increasing the production of reproduction releasing hormones from the hypothalamus. This was supported by the findings of Gong, *et al.* (2002), who reported that the high insulin including diet reduced the PPI and increased the conception rate in dairy cows.

- **Corticosteroids**

During the final stages of gestation the plasma concentration of cortisol increases. Due to the rapid growth of the fetus and increasing the demand to the metabolic agents this stimulates the placenta to produce  $\text{PGF}_2\alpha$  which stimulates the fetal hypothalamus-pituitary and adrenal axis leading to increase concentration of cortisol (Hafez, 1993). The cortisol stimulates the placenta to convert progesterone to oestrogen. The elevated level of oestrogen stimulates secretion of  $\text{PGF}_2\alpha$  from uterine wall and enhance developing of oxytocin receptors.

#### **1.4.3.7 Milking:-**

Recent studies carried on cows with high genetic merits kept under a grass based production system indicated that they had long interval to first service (Buckley, *et al.* 2000 a and b). High genetic merit is associated with high milk yield. The same above mentioned authors suggested a poor reproductive performance in high milk producing cows, as reflected by a poor conception rate. The high milk yield is reported to have a negative effect on recrudescence of oestrous postpartum (Harrison, *et al.* 1990).

#### **1.4.3.8 Heat stress:-**

The climatic factors and husbandry practices are important constraints affecting dairy production (Ageeb and Hillers, 1991). The heat stress is the major environmental factor affecting fertility of dairy cows during hot season (Abilay, *et al.* 1975). The Same authors reported that the heat stress reduces plasma oestradiol during pro-oestrus, and affects the length and intensity of oestrous behavior. This explain the smaller proportion of cows detected in oestrus during Summer season (Gwazdauskas, *et al.*, 1981). Many alternatives have been made to reduce the effects of heat stress on dairy cows through the use of shade, fans, air conditioning and sprinklers, (Thatcher, 1974; Wise, *et al.*, 1988; Johnson, 1991). The heat stress reduces the feed intake, milk production and increase mortality rates. The higher temperature in the tropics is inimical to sustainable dairy production using Bos Taurus, (Vaccaro, 1974; Ansell, 1985; Eisen, 1989). The importation of temperate breeds for cross breeding in Sudan started in 1925. Tendency to keep purebred Holstein- Friesian cow for intensive dairy system has developed in recent years (Osman, 1984).

#### **1.4.3.9 Uterine infections**

Uterine infections delay the initiation of folliculogenesis (Peters and Bosu, 1988). The uterine infections cause suppression of the rate of follicular growth in dairy cows during early puerum by inhibiting LH release. The same authors found that, the inhibition was due to endotoxins produced by gram-negative bacteria in the postpartum period.

Postpartum uterine infections occur commonly in the cows due to retention of fetal membranes and dystocia (Hafez, 1993). Placental retention accompanied with metritis was known to delay uterine involution (Arthur, *et*

*al.*, 1998). Pyometra in the cow was reported to develop when ovulation occurs within 3 weeks after calving with presence of large numbers of pathogenic bacteria in the uterus (Olson, *et al.*, 1984).

#### **1.4.10 Management:-**

The poor standard of herd management will lead to poor detection rates of oestrus (Arthur, *et al.* 1998). Therefore a well trained staff for herd management of dairy farms is needed to identify oestrus correctly. The importance of trained staff for herd management was also stated by Esselmont and Kossaibati (2000) who found that the percentage of cows identified in oestrus by a well trained herd staff ranged between 82-97% where percentage of cows in oestrus detected by untrained staff was 67%. There are some constrains like over crowding and muddy floors which will reduce the accuracy of oestrous detection because cows that are not in oestrus will not be able to escape away from other mounting cows (Arthur, *et al.* 1998).

#### **1.4.11 Breed:**

The breed is considered to be as the main constrain affecting the reproductive performance of the dairy cows therefore different breeds had a different length of PPI (Ruiz, *et al.* 1992). The genetic influences of the breed can be demonstrated by the short period for the return of ovarian function postpartum in the dairy breeds 10-45 days compared with beef breeds 36-70 days (Arthur, *et al.*, 1998). Crosses between Bosindicus and temperate cattle are known to have a relative longer PPI compared to crosses among temperate cattle (Estrada, *et al.* 1992).

### **1.5 Biological Uses of Gonadotropin – releasing hormone (GnRH)**

GnRH is a decapeptide hormone synthesized in cell bodies of neurosecretory neurons located in the mediobasal hypothalamus (Arthur, 1998). It stimulates the pituitary gland to release LH and FSH (Thatcher, *et al.*, 1991). The GnRH effects can be indirect through its effect on pituitary to release LH and FSH (Chenault, *et al.*, 1990) or direct on the reproductive tissues. Two potential gonadotropin responsive tissues within the ovary are the corpus luteum (CL) and the follicle.

Thatcher, *et al.* (1993) reported that the GnRH will not have direct stimulatory or inhibitory effects on CL to increase progesterone production. Thus, potential GnRH effects must be mediated indirectly via alterations in endogenous gonadotropin secretion or directly via non receptor mediated mechanism. The GnRH and its agonist analogues act on ovarian follicular development and CL function indirectly via the induced release of pituitary LH and FSH (Hafez, 1993). Administration of GnRH increases LH and FSH in the peripheral circulation within 2-4 hours (Chenault *et al.*, 1990; Stevenson *et al.*, 1993). These gonadotropins act directly by binding to their respective receptors on follicular and luteal cells (Mac Millan, and Thatcher, 1991). The GnRH causes inhibition of oestrus and induces ovulation of large follicles. The induction of ovulation of the large follicle is associated with decrease in oestradiol concentration ( $E_2$ ) in the peripheral circulation. Ovulation does not occur in all cases after GnRH treatment because the ability of a follicle to ovulate depends on its developmental stage. The concentration of progesterone decreases  $<1$  n mole/ml and the large follicle produces oestradiol that is involved in triggering oestrus via its positive feedback on hypothalamus- hypophysis center (Hafez, 1993). This stimulates the release of the GnRH and induces LH pulses and the LH surge which is

responsible for ovulation of the large follicle. The delay of oestrus following treatment with a GnRH agonist is due to functional alterations of large follicle present at the time of treatment (Twagiramungu, *et al.* 1995).

The GnRH is a potential drug that is used widely in the management of postpartum reproductive efficiency in dairy cows (Fernandez, *et al.*, 1978). Administration of The GnRH eliminates the large follicles by ovulation or atresia and initiates new follicular waves within 3-4 days after treatment at any stage of oestrous cycle (Twagiramungu, *et al.*, 1995). It is also found that administration of the GnRH to the dairy cows postpartum, reduces the PPI (Thatcher, *et al.* 1993). It is found that the pregnancy rate reached 53.7% for the cows injected with GnRH on the day of parturition (Mialot, *et al.* 1999). The same authors reported that, cows given the GnRH on the day of parturition showed oestrus on day 10. Elzubeir and Elsheikh (2004) reported that, cross bred dairy cows (Kenana  $\times$  Friesian ) when received an injection of GnRH during the 1<sup>st</sup> 2<sup>nd</sup> or the 3<sup>rd</sup> week postpartum has a similar number of services per conception ( $1.00 \pm 0.00$ ) which were lower than that of the control ( $1.40 \pm 0.20$ ). Moreover the same authors reported that the GnRH treated cows have a shorter days open and calving interval than untreated cows.

The resumption of ovarian activity and oestrus early in the postpartum period can be enhanced by a single injection of GnRH given at different times after calving (Schams, *et al.*, 1973; Britt, *et al.*, 1977; Irvin, *et al.* 1981; Boiti, *et al.*, 1982; Ball and Lamming, 1983).

## **1.6 Calving Interval (CI)**

It is the period between two consecutive calving. It was reported that the CI of zebu cattle in North Sudan is 428 days. While that of temperate cows namely Friesian bred in Sudan was a round 360 days (Bashir, 1990). A calving interval (CI) close to 12 month is generally regarded as the most profitable length of the CI in dairy production (Zeddies, 1982; Strandberg and Oltenacu, 1989). This assumption is mainly based on economical calculations on milk production, feeding and housing. The CI is an objective trait of economic importance (Olori, and Galesloot, 2000). Also the CI can be considered as a good indicator of cow fertility (Campos, *et al.* 1997; Pryce, *et al.* 1998).

The CI is influenced by the same factors that influence the PPI



## **Materials and Methods**

### **2.1 Study area**

This study was done in Nyala City which is the capital of South Darfur State of Western Sudan, it lies within the area between latitude  $12^{\circ} 03' 25''$  N and longitude  $24^{\circ} 53' 09''$  E, with 671m above the sea.

### **2.2 Climate:-**

South Darfur State has three different climates. Semi desert climate with average rain fall of about 100-300mm/year. Poor Savanna climate with average rain fall of about 300-600mm/year. Rich Savanna climate with average rain fall of about 600-900mm/ year.

There are three distinct seasons through out the year. The hot dry Summer season from March to June, rainy season from July to October, and Winter season from November to February.

### **2.3 Temperature measurement:-**

The daily temperatures were obtained from Nyala Meteorological Department (appendix .4.)

The mean temperature during Winter is (32.3°C maximum, 13°C minimum), during Summer is (38°C maximum, 21.4°C minimum) and during Autumn is (34.3°C maximum, 17.9°C minimum). (See appendix 4)

**Table .2.**

**The average of annual rain fall /per/ mm in Nyala station from 1999-2005**

<b>The year</b>	1999	2000	2001	2002	2003	2004	2005
<b>Nyala Station</b>	480.6	397.4	286.2	311.1	552.4	268.00	494.00

See appendix .1.

## **2.4 Animals**

In this study three local dairy breeds of cows native to South Darfur State were employed (n = 193 cows). These breeds were, Kenana breed, n = 72 Fellata breed n = 60 and cross bred (Kenana × Friesian) n = 61. The BCS of these cows vary between 2.5-4, according to the scale described by Wildman, *et al.*, (1982). Their parity range was between 1 to 5.

### **2.4.1 Management and feedin**

The Kenana and Fellata cows were kept under semiclosed system of management. During the day the cows were allowed to graze on the surroundings from 10:00 am and flocked back at 4: 00 pm. They were fed roughtages and concentrate mixture which is composed of Sorghum (Sorghum Vulgare vr Fetarita) 35%, ground nut cake 30% wheat bran 33%

and Sodium chloride 2%. The Kenana and Fellata were fed 3-4 kg concentrate once a day in the afternoon during the milking period.

The cross-bred cows were kept under closed system of management. They were fed dairy concentrate mixture at a rate of 5kg / cow in the morning and 5kg/cow in the afternoon. The concentrate mixture is composed of 38% Sorghum, (Sorghum Vulgare vr Fetarita) 30% groundnut cake 30% wheat bran and 2% sodium chloride. They were fed green fodder Abu70 (Sorghum, bicolor), Barseem (Alfa Alfa) and roughages.

After parturition, the new-born calves were left with their dams for 7 days and thereafter they were left to suckle their mothers twice a day in the morning and in the afternoon.

#### **2.4.2 Housing :**

The animals were kept in an open shaded yards. The open shaded yards were constructed mainly from local materials. The pens were made from wooden bars and the roof was from locally made wooden sheets. The floor was covered with sand. The pregnant animals were kept in separate yards.

#### **2.4.3 Milking:**

The average daily milk production during the first 90 days postpartum was 3-5 lb/day for Fellata breed, 10-12 lb/day for Kenana breed and 20-30 lb for cross-bred cows. After parturition the dams were left to suckle their calves twice daily till the end of lactation period which ranged between 6-7 months. Milking was done manually by the herd men.

#### **2.4.4 Heat detection**

All the cows in the herds were checked for oestrous signs by visual observations by well trained herdsmen three times a day postpartum, early in the morning at 7:00 am, in the mid day at 12:00 and at 6:00pm for at least 30 minutes. The cow was recorded in heat when it becomes restless, licks the perineum of other cows, jumps on other cows, allow other cows or bull to mount her, it bellows and there is a transparent clear mucous drops from her vulvae. The cow was considered, in a full response when it stands to be mounted by the bull and mating was completed. (Arthur, *et al.* 1998).

#### **2.4.5 Health control.**

Routine vaccination against the common major diseases Haemorrhagic Septicemia (HS), Black quarter (BQ), Anthrax and Contagious Bovine Pleuropneumonia (CBPP), was done once yearly. Routine testing for brucellosis was done yearly under the control of the Regional Veterinary Research Lab. Routine mastitis testing was done for the suspected cows. The udder was washed with water and after milking the teats of the udder were emerged in a suitable antiseptic solution like (potassium permanganate 1:1000). During the last week prepartum the pregnant cows were injected with streptomycin 20% to prevent postpartum mastitis. They were also treated with local intramammary antibiotic to reduce the incidence of mastitis during the dry period. The commonly used antibiotic in the farms as treatment of respiratory infections was oxytetracycline at a dose rate of 50 mg/ 5-10 kg. b.w. (Limoxin -50

Interchemec Holland). The cows were drenched Anizole-100 suspension 10 mg/ kg b.w. (Anglian Nutrition Products Company United Kingdom) the shades were routinely sprayed with acaricidal drugs.

## **2.5 Experimental design:-**

### **2.5.1 Experiment -I-**

This experiment was a simple factorial design to study the breed effect on the PPI of three Sudanese dairy breeds of cattle native to South Darfur State. The length of the PPI was studied by visual observations. A total of 59 dairy cows calved between January 2003-April 2004 were selected according to the scale of the BCS after calving. The cows with BCS below 2.5 were excluded. The general range of the BCS for the cows used in this experiment was between 2.5 -4 with parity between 1-5. The cows were grouped into three groups according to their breeds, Kenana (n = 21), Fellata (n = 17) and cross-bred (Kenana  $\times$  Friesian) (n = 21) cows. The length of the PPI was calculated from the day of parturition until the appearance of the first signs of the first oestrus postpartum. Furthermore, the influences of BCS and parity on the PPI were investigated.

### **2.5.2 Experiment -II-**

This experiment was a 3 $\times$ 2 factorial design to study the possibility of reducing the length of the PPI of the three local breeds of dairy cattle Kenana, Fellata and cross-bred native to south Darfur State treated with gonadotrophic releasing hormone (GnRH) on day 21 postpartum. A total of (n = 66) dairy cows were selected for this experiment. The scale of the BCS

for the cows used in this experiment ranged between 2.5-4 after calving, the parity ranged between 1-5. The cow with BCS less than 2.5 were excluded. The cows were grouped into two groups, group I was the treatment group (n = 33) and group II was the control (n = 33). Group I was subgrouped into three groups according to their breed, Kenana (n=11) Fellata (n = 11) and cross-bred (Kenana  $\times$  Friesian) (n = 11) cows. Every cow from group I was intramuscularly injected with 250 $\mu$ g of synthetic GnRH (Fertagyl, Intervet International B.V. Boxmeer, Hollanda) on day 21 after calving (Elzubeir and Elshiekh, 2004). Group II was untreated to serve as control. The length of the PPI was calculated from the day of parturition till the appearance of the signs of the first oestrus postpartum.

### **2.5.3 Experiment -III-**

This experiment was a 3 $\times$ 3 factorial design to study the effect of season of calving on the length of the PPI of the dairy cows native to South Darfur State. The total number of cows used in this experiment was 68 dairy cows, they were grouped into three groups according to the breed, Kenana (n = 29) cow, Fellata (n = 21) cow and cross bred (Kenana  $\times$  Friesian) (n = 28) cow. Each group was further sub grouped according to the season of parturition. The cows that calved in Summer season were (n = 25) dairy cows, Kenana (n = 7), Fellata (n = 4) and cross bred (n = 14) cows. The cows that calved in Winter season were (n = 31) dairy cows, Kenana (n = 13), Fellata (n = 11) and cross-bred (n = 7) cows. The cows that calved in Autumn season were (n = 22) cows, Kenana (n = 9), Fellata (n = 6) and cross-bred (n = 7) cows. The BCS of the cows used in this experiment was between 2.5-4 after parturition. The length of the PPI was calculated from the day of parturition until the appearance of the signs of the first oestrus postpartum.

## **2.6 Statistical analysis**

Data are means  $\pm$  SE. The data were subjected to ANOVA followed by Fisher's protected least significant difference. Differences at probability of  $P < 0.05$  were considered statistically significant. The relation between the BCS and the PPI, and parity and PPI were blotted using the regression blot.

## **Results**

### **3.1 Experiment .I.**

#### **3.1.1 The effect of the breed on the length of the PPI.**


As shown in **Fig. 1.** breed significantly ( $P < 0.001$ ) affects the length of the PPI. The cross-bred dairy cows had a shorter PPI ( $122.00 \pm 14.9$  days) compared to the kenana breed ( $286.9 \pm 35.3$  days) and the Fellata breed ( $246.26 \pm 26.6$  days). No significant difference ( $P > 0.05$ ) was observed between the kenana and the Fellata breed.

#### **3.1.2 The relation between the BCS and the PPI.**

As shown in **Fig. 2.** there was a slight negative relation between the BCS and the length of the PPI ( $R=0.2$ ). When the BCS increases the length of the PPI decreases.


#### **3.1.3 The relation between the parity and the PPI.**

As shown in **Fig. 3.** parity has no influence ( $R=0.005$ ) on the PPI of the different breeds of cows, however when the number of parities increases the PPI decreases.






**Fig. 1.** Postpartum interval in three different dairy breeds bred in South Darfur state. <sup>a, b</sup> Values differed at  $p < 0.001$ .



**Fig. 2.** The relation between the BCS and the length of the PPI of the local breeds Kenana, Fellata and cross-bred dairy cows.



$$Y = 245.303 - 9.714 * X; R^2 = .005$$

**Fig. 3.** The relation between parity and the length of the PPI of the local breeds Kenana, Fellata and cross-bred dairy cows.


### 3.2 Experiment -II-

### **The effect of treatment with GnRH on day 21 postpartum on the length of the PPI.**


The results of this experiment showed that there was no interaction ( $P>0.05$ ) between the breed and treatment. Treatment with GnRH significantly ( $P<0.001$ ) reduces the length of the PPI in all breeds of cows (Kenana, Fellata and Cross bred). The length of the PPI of the Kenana cows treated with GnRH on day 21 postpartum was  $79.5 \pm 8.4$  days and the control Kenana cows was  $276.5 \pm 54.4$  days. (**Fig. 4**). The PPI of Fellata cows treated with GnRH on day 21 postpartum was  $65.9 \pm 3.2$  days while that of the control Fellata cows was  $266.4 \pm 38.7$  days. (**Fig. 5**)

**Fig. 6.** Explains that the PPI of the treated cross-bred cows was  $74.2 \pm 2.8$  days, while that of the control cross-bred cows was  $151.9 \pm 18.5$  days.


Also the results showed no significant ( $P>0.05$ ) difference among the treated cows of the different breeds.



**Fig.4.** The effect of GnRH treatment on day 21 postpartum on the length of PPI of Kenana cows. <sup>a, b</sup> Values differed significantly at  $p < 0.001$ .



**Fig.5.** The effect of GnRH treatment on day 21 postpartum on the length of PPI of Fellata cows. <sup>a, b</sup> Values differed significantly at  $p < 0.0001$ .



**Fig. 6.** The effect of GnRH treatment on day 21 postpartum on the length of PPI of cross-bred cows. <sup>a, b</sup> Values differed significantly  $p < 0.001$ .

### **3.3Experiment -III-**


#### **Effect of the season on the PPI**

As shown in **Fig. 7**. the season significantly influences the PPI of cross-bred cows ( $P<0.001$ ) and it has slight effect on the PPI of Kenana, and Fellata cows. The cross bred cows in Summer expressed a significantly ( $P<0.01$ ) short PPI compared to Winter and Autumn.

The length of the PPI of Kenana cows during Winter, Summer and Autumn were,  $276.4 \pm 46.9$  days,  $214.60.0$  days,  $223.5 \pm 31.6$  days, respectively .

The length of the PPI of Fellata cows during Winter, Summer and Autumn were,  $286.5 \pm 38.7$  days,  $200.8 \pm 30.4$  days and  $223.2 \pm 22.9$  days respectively. The length of the PPI of cross-bred cows during Winter, Summer and Autumn were,  $153.1 \pm 17.5$  days;  $46.14 \pm 5.1$  days and  $160.5 \pm 29$  days respectively.





**Fig. 7.** The effect of the season on the length of the PPI of the three breeds, Kenana, Fellata and cross bred dairy cows.  
a, b, c Values differed significantly at ( $P>0.01$ ).

## **Chapter -4-**

### **Discussion**

In the present study it was found that the cross-bred dairy cows had a shorter PPI than the pure local breeds, namely Kenana and Fellata cows native to South Darfur State. It has now become evident that the reproductive performance has been decreasing in the high yielding cows. A large number of studies have indicated that milk yield and genetic merits for milk production are negatively associated with fertility traits (Campos, *et al.* 1997.; Hoekstra, *et al.* 1984; Pryce, *et al.* 1997). Other studies have found that the reproductive function during the early postpartum period is reduced in animals with high genetic merit or high milk yield (Beam and Butler, 1997; Snijers, *et al.* 2001). Although it is well known that the temperate dairy cows produce large amount of milk compared to the zebu cows, the length of their PPI is shorter than that of zebu cows. In the current study the cross-bred dairy cows used were produced by crossing of Friesian with Kenana to increase the genetic merits for milk production. Results from the present studies have provided evidence that cross-bred dairy cows having Friesian blood have a shorter PPI compared to the Bous indicus cows namely Kenana and Fellata dairy cows native to Sudan. Although the two

pure native breeds of cows used in the current study produce less milk than the cross-bred cows used. Their reproductive function during early postpartum is slow than that of the cross-bred cows. It seems that the negative relation between increasing genetic merit and the reproductive function during early postpartum in the previous studies takes place only when selection for improvement is carried out within the same breed. However, results from the present study shown that up grading of zebu cows with Friesian to increase their genetic merit for milk yield, reduced their PPI and did not increase it. This finding indicated that there exists a genetic factor in temperate cows not related to milk yield that makes them have a shorter PPI.

It has been proposed that the reproductive performance is influenced by the negative energy balance during early lactation (Staple, *et al.*1990; Beam and Butler, 1997). Owing to the difference between dietary energy intake and requirement for milk production, dairy cows experience a period of negative energy balance during early lactation which results in mobilization of body fat reserve (Peters and Riley, 1982). The extent for negative energy balance is normally related to the milk yield (Butler and Smith, 1989; Beam and Butler, 1997). Several studies have shown that the negative energy balance status can significantly influence functional characteristic of ovarian follicular development during early postpartum period and therefore affects the resumption of oestrous cycles, interval from calving to the first ovulation and subsequent fertility (Staples, *et al.*; 1990; Lucy, *et al.* 1992; Beam and Butler, 1997; Kruip, *et al.* 1998). The negative energy balance that leads to mobilization of body fat reserve is associated with changes in BCS score. The cows that loose more fat will have a less

BCS. The results of the current study showed that there is a slight negative relation between the length of the PPI and the BCS. That is to say the cows that their BCS decreases during early postpartum will express a longer PPI compared to the cows that their BCS remains stable. However, other studies have reported that the PPI to first oestrus is not related to energy balance of the cow (Villa-Godoy, *et al.* 1988.; Snijders, *et al.* 2001) and that the BCS from calving to first service are not related to conception in high genetic merit cows (Snijders, *et al.* 2001). The difference between these studies and the present study was probably due to differences in metabolic hormone profile which is known to influence the resumption of oestrous cycle during early postpartum.

Postpartum ovulation occurred earlier in cows that had calved more than twice compared to cows that calved twice or less (Eduvie, 1985). The primiparous cows took longer time to return to first oestrus postpartum than did pluriparous cows (Mukasa, *et al.* 1991). The shortest time taken for occurrence of first oestrus was reported to be at the 3<sup>rd</sup> - 4<sup>th</sup> calving, the moderate period was at the 5<sup>th</sup> - 6<sup>th</sup> calving and the longest time taken was at 9<sup>th</sup> - 10<sup>th</sup> calving (Elhag, 2003). The findings of the present study indicated that the parity has no effect on the length of the PPI of dairy cows native to South Darfur State, these findings disagree with the above mentioned studies. This difference may be due to the range of parity used for the present study which was between 1<sup>st</sup> to 5<sup>th</sup> calving, where that of the above studies may extend to 10<sup>th</sup> calving. Additionally, the cattle owners in South Darfur State used to breed their heifers at the age of 3 years or above, consequently the age at 1<sup>st</sup> calving of the cows native to South Darfur State

will be different from those cows used in the above studies. The age at 1<sup>st</sup> calving of the cows in the above mentioned studies was about 2.5 years.

Administration of GnRH during early postpartum causes an acute increase in gonadotropins level (FSH and LH) that stimulate folliculogenesis, induce ovulation of the selected large follicle and increase oestrogen concentration in the plasma (Hafez, 1993). Oestrogen is known to have a bactericidal effect that reduces infection during postpartum period resulting in a faster uterine involution and a reduced PPI (Thatcher, *et al.* 1993). A single injection of GnRH during the 1<sup>st</sup>, the 2<sup>nd</sup> or 3<sup>rd</sup> week postpartum reduced the time taken for recrudescence of the postpartum first oestrus in cross-bred (Friesian × Kenana), Sudanese dairy cows (Elzubeir and Elsheikh, 2004). Early postpartum administration of GnRH initiated the postpartum ovarian cyclic activity and reduced the PPI (Yavas and Walton, 2000). The GnRH and its agonist analogues act on ovarian follicular development and CL function indirectly via the induced release of pituitary LH and FSH that stimulates ovarian activity leading to reduction of the length of the PPI (Conn and Crowley, 1991). In the present study injection of GnRH on day 21 postpartum reduced the length of the PPI. This result agrees with the finding of the above mentioned studies. Contrary (Thatcher, *et al.*; 1993; Foote and Rick, 1999) reported that injection of GnRH during early postpartum has no effect on the reproductive efficiency on the time elapsed from parturition to the occurrence of first oestrus. These discrepancies are probably due to differences in the time of GnRH administration which was carried on day 21 postpartum in this study and in

the above studies, the time of administration of GnRH was before the 3<sup>rd</sup> week postpartum. Also the effect of breeds employed could not be ignored.

Seasonal changes in temperature and photoperiod are known to influence the timing of recrudescence of the first oestrus postpartum (Hafez, 1993). Along postpartum period was reported in cows that delivered in Winter and early spring than those delivered in Autumn and Summer (Peter and Riley 1982). The same authors reported a long PPI in cows calved in Spring compared to cows calved in Autumn. Furthermore, Peter and Riley reported that expression of the first postpartum oestrus during Winter is lower than it is expression in Summer. In the current study the pure local cows namely Kenana and Fellata cows showed similar PPI among all seasons. However the cross-bred dairy cows in the current study expressed a shorter PPI during Summer compared to Winter and Autumn. The findings of the cross-bred cows agree with the above mentioned studies. Minor discrepancies may occur between the findings of this study and the above mentioned studies because in South Darfur State there are only three seasons during the year of which Winter is very short and not well pronounced and there is no Spring season. These difference (namely breeds and seasons differences) may also act individually or cooperatively to make the length of the PPI of the pure local cows looks similar in all seasons of the year.

## **Conclusion**

From the present study it concluded that:

- ➔ Kenana cows bred in South Darfur State have a mean PPI of  $286.9 \pm 35.3$  days.
- ➔ Fellata cows have a mean PPI of  $246.3 \pm 26.6$  days.
- ➔ Cross-bred dairy cows native to South Darfur State have a mean PPI of  $122.0 \pm 14.9$  days.
- ➔ Local dairy cows having foreign blood have a shorter PPI than pure local breeds.
- ➔ The BCS has a slight negative relation with the length of the PPI. When the BCS increases the length of the PPI decreases.

- ➔ Parity up to 5 has no effect on the length of PPI of Kenana, Fellata and cross bred dairy cows native to South Darfur State.
- ➔ The season has no effect on the PPI of pure local cows namely Kenana and Fellata.
- ➔ The PPI of cross-bred dairy cows native to South Darfur State is affected only by Summer season. While no effect of Autumn and Winter on PPI of the cross- bred dairy cows was recognized.
- ➔ Increasing the genetic merits for milk yield by crossing the local breed of cows with foreign breeds reduces the length of PPI.
- ➔ Injection of GnRH on day 21 postpartum reduces the length of PPI of all breeds.



### **Recommendations:**

- 1- To obtain dairy cows that have a short PPI and capable for producing a calf per year, the local dairy cows native to South Darfur State should be up graded with foreign breeds of cows.
- 2- The BCS of the cows during pregnancy should be above three to avoid the loss in the body weight during early postpartum that leads to a longer PPI.
- 3- A program of administration of GnRH after the 3<sup>ed</sup> week postpartum is recommended to reduce the length of the PPI.

## Reference

- Abilay, T.A.; Johnson, H.D. and Madan, M. (1975). Influence of environmental heat on peripheral plasma progesterone and cortisol during the bovine estrous cycle. *J. Dairy Sci.* 58:1836-1840.
- Acosta, B.G.K.; Tarnavsky, T.E.; Platt, D.L.; Hammernick, J.L.; Brown, H.M.; Schoenemann, H.M. and Reevels, J.J. (1983). Nursing enhances the negative effect of oestrogen on LH release in the cow. *J. Anim. Sci.* 57: 1530-1542.
- Ageeb, A. and Hillers., J.K. (1991). Production and reproduction characteristics of Butana and Kenana cattle of the Sudan. *World Anim. Rev.* 2: 49-56.
- Ansell, R.H. (1985). Cattle breeding in the tropics. *World Anim. Review.* 20:73-79.
- Arthur, G.H.; Noakes, D.E.; Harold, P. and Parkison, T.J. (1998). Infertility in the cow. In: *Vet. Reproduction and Obstetrics* (7th ed.) W.B. Saunders London Publishers, PP 345-388.

Ball, P.J.H. and Lamming, G.E. (1983). Diagnosis of ovarian acyclicity in lactating dairy cows and evaluation of treatment with gonadotrophin-releasing hormone or progesterone releasing intravaginal device. Br. Vet. J. 139: 522-527.

Bashir, I.E. (1990). *Reproductive performance of Holstein Frisian Dairy Cows with indirect different cross-bred Grades in the Sudan*. M. Sc. Thesis. Faculty of Anim. Prod. U of K.

Beam, S.W. and Butler, W.R. (1997). Energy balance and ovarian follicle development prior to the first ovulation postpartum in dairy cows receiving three levels of dietary fat. Biol. Reprod. 56: 133-142.

Benmard, M. and Stevenson, J.S. (1986). GnRH and PGF2 $\alpha$  for postpartum dairy cow's oestrous, ovulation and fertility traits. J. Dairy Sci. 69: 800-811.

Boiti, C.; Biphelli, V.; Simontacchi, C.; Olivieri, O.; Caggioni, C. and Micale, R. (1982). Postpartum progesterone profiles and reproductive performance of dairy cows treated with GnRH or progesterone. In: *Factors influencing Fertility in the postpartum cows*. H. Karag and E. Schallenberger, (ed) Martinus, Nijhoff, the Hague, PP. 544-547.

Britt, J.H.; Harrison, D.S. and Morrow, D.A. (1977). Frequency of ovarian cycles: reason for culling and fertility in Holstein Friesian cows given GnRH at two weeks after parturition. *Anim. J. Vet. Res.* 38: 749-751.

Buckley, F.; Dillon, P.; Cross, S.; Flynn, F. and Rath, M. (2000a). The performance of Holstein Friesian dairy cows of high and medium genetic merit for milk production on grass based feeding systems. *Livest. Prod. Sci.* 64: 107-119.

Buckley, F.; Dillon, P.; Rath, M. and Veerkamp, R.F. (2000b). The relationship between genetic merit for Yield and Live weight, condition score and energy balance of spring calving Holstein Friesian dairy cows on grass based system of milk production. *J. Dairy Sci.* 83: 1878-1886.

Butler, W.R.; Everett, R.W. and Coppick, C.E. (1981). The relationships between energy balance, milk production and ovulation in postpartum Holstein cows. *J. Anim. Sci.* 53: 742-749.

Butler, W.R. and Smith, R.D. (1989). Interrelationships between energy balance and postpartum reproductive function in dairy cattle. *J. Dairy Sci.* 72: 767-783.

Campos, M.S.; Wilcox, C.L.; Beceril, C.M and Dig. A. (1997). Genetic parameters for yield and reproductive traits in Holstein and Jersey cattle in Florida. J. Dairy Sci. 77: 867-873.

Central Bureau for Statistic, (2004). Population data sheet for Sudan by state CBS-Khartoum. Sudan.

Chenault, J.R.; kratzer, D.D; Rzepkowski, R. A. and Goodwin, M. C (1990). L H and FSH response of Holstine heifers to fertirelin Acetate, Gonadorelin and buserlin. Theriogenology, 34: 81 - 98.

Conn, P.M. and Crowley W.F (1991). Review Article: Gonado trophin releasing hormone and its analogues. New Eng. J Med. 324: 93 - 103 .

Coppock, C.E.; Noller, C.H. and Walfe, S.A. (1974). Eddecty of forage concentrate ratio in complete feeds ad libitum on energy. J.Dairy Sci. 57: 1371-1380.

Dimmick, M.A.; Gimenez, T. and Spitzer, J.C. (1991). Ovarian endocrine activity and postpartum interval in beef cows. Anim. Reprod. Sci. 24: 173-178.

Eduvie, L.O. (1985). Factors affecting postpartum ovarian activity and uterine involution in Zebu cattle indigenous to Nigeria. Anim. Reprod. Sci. 8: 123-128.

Eisen, E.J. (1989). Proceeding of Symposium on utilization of Animal Genetic Resources in latin America, Brazil. J. of Genetics, 12 (supplement): 13-26.

Elhag, M.A.S.; (2003). *Factors influencing postpartal reproductive traits of cross-bred dairy cows in the Sudan*. M. Sc. Thesis. Faculty of Veterinary Medicine, U of K.

ELkeraby, F.E. and Abu-Ela, M.B. (1982). A study of some non genetical factors affecting the postpartum reproductive performance in Friesian cows. Trop. Anim. Prod. 7: 307-317.

Elsheikh, A.S. and Ahmed, F.O. (2005). Backing up postpartum dairy cows with PGF<sub>2</sub> $\alpha$ . J. Anim.Vet. Advan. 4 (5): 506-509.

Elzubeir, F.O. and Elsheikh, A.S. (2004). Reproduction performance of cross-bred Sudanese dairy cows treated with GnRH during early postpartum. J. Anim. And Vet Advances. 3 (5): 329-334.

Elzubeir, F.O.A. (2003). *Reproductive performance of cross-bred Sudanese dairy cows treated with PGF<sub>2</sub> $\alpha$  or GnRH during early*

*postpartum period*. M. Sc. Thesis. Faculty of Veterinary Medicine, U of K.

Esslemont, R.J. and Kossaibati, M.A. (2000). The use of database to manage fertility. J. Anim. Reprod. Sci. 61: 725-741.

Estrada, M.F.; Gonzales, S.C. and Villar, A. (1992). Effect of milk yield on reproductive performance and efficiency in primiparous cows. Anim. Breed Abst. 60 (4): 284-296.

Fernandez, L.G.; Thatcher, W.W.; Wilcox, C.J. and Call, E.P. (1978). LH Release in responses to GnRH during the postpartum period of dairy cows. J. Anim. Sci. 46: 443-448.

Foote, R.H. and Riek, P.M. (1999). GnRH improves reproductive performance of dairy cows with slow involution of the reproductive tract. J. Anim. Sci. 77 (1): 12-16.

Galina, C.S. and Arthur, G.H. (1989). Review of cattle production in tropics. Part 2. Anim. Breeding Abst. 57: 679-686.

Ginther, O.J; Kot , K ; Kulick , L.J. ; Martins and Witbank, M.C. (1996). Relationships between FSH and ovarian follicular waves during the Last six months of pregnancy in cattle. J. Reprod. Fertil. 108: 271- 282.

Gong, J.G. Lee, W.J; Garnsworthy, P.C. and Webb, R. (2002). Effect of dietary concentrations during the early postpartum period on reproductive function in dairy cows. J. Reprod. Fertil. 123: 419-227.

Gong, J.G.; Bramely, T.A.; Gutierrez, C.G.; Peters, A.R. and Webb, R. (1996). Effect of chronic agonist on peripheral circulation. J. Reprod. Fertil. 105: 263-270.

Gwazduskas, F.C.; Thatcher, W.W.; Kiddy, C.A.; Paape, M.J. and Wilcox, C.J. (1981). Hormonal patterns during that stress following PGF<sub>2</sub> $\alpha$  that induced luteal regression in heifers. Theriogenology, 16: 217-285.

Hafez, E.S.E. (1993). Reproductive cycle. In: *Reproduction in Farm Animals 6th Edition* Hafez. E.S.E.(ed) A Lea and Fibiger Kiawah Island, South Carolina, USA pp. 230-260.

Harrison, R.O.; Ford, S.P; Young, V.W. and Freemans (1990). Increased milk production versus reproductive and energy status of higher producing dairy cows. J. Dairy Sci. 73: 2749-2758.



Hoekstra, J.A.W.; Vander, J.H.J. and Quweltjes, W. (1984). Genetic and phenotypic parameters for milk production and fertility traits in up graded dairy cattle. *Livest. Prod.Sci.* 40: 225-232.

Irvin, H.J.; Zaied, A.A.; Day, B.N.; Garverick, H.A. (1981). GnRH induced postpartum an estradiol 17B concentration on the release of LH following administration of GnRH. *Theriogenology*, 15: 443-456.

Johnson, H.D, 1991. The lactating cow in the various ecosystems, environmental effects on its productivity, FAD Animal production and health paper, 9: 9 - 21.

Kindahl, H.L; Edqvist, E. ; Larson , K. and Malmquist, A. (1982) . Veterinary medicine and animal science in: *Current Tropic in Veterinary Medicine*. Krag , H; schallenperger , E.D. S (ed). Martinus, Nighoff. The Hague, pub. 20:173- 196.

Kruip, T.A.M.; Meyer, G.A.L.; Rukkwamsuk, T. and Wensingt (1998). Effects of feed in the dry period on fertility of dairy cows postpartum. *Reprod. in Dom. Anim.* 33: 165-168.

Kovacevic, R.; Krasmanovic, L.; Maric, D.; Perkucin, R. and Veselinovic, S. (1988). Investigation of hormone profiles in dairy cattle with

a view controlling fertility and increasing reproductive efficiency. Proceeding of final research coordination meeting organized by the joint FAO/ IAEA and cooperation Italiana Allo Sviluppo, held in Rabbat from 23-27 March 1987. Vienna (Austria). IAEA. 379: 255-274.

Lamming, G.E; Wathe, D.C and Peters, A.R (1981). Endocrine patterns of the postpartum cow. J . Reprod. Fertil. 30: 155- 162.

Lamming, G.E.; Peters, A. R.; Rilery, G.M. and Fisher, M.W. (1982). Endocrine regulation of postpartum function.J. Anim. Sci. 20: 143-160.

Lindel, J.O.; Kindahl, H; Jansson, L. and Edqvist, L.E. (1982). Postpartum release of PG F<sub>2</sub> $\alpha$  and uterine involution in the cows. Theriogenology, 17: 237- 245.

Lucy, M.C.; Staples, C.R.; Thatcher, W.W.; Erickson, P.S.; Cleale, R.M.; Firkins, J.L.; Clark, J.H.; Murphy, M.R and Brodie, B.O. (1992). Influence of diet composition, dry matter intake, milk production and energy balance on time of postpartum ovulation and fertility in dairy cows Anim. Prod. 54:323-331.

MacMillan, K.L. and Thatcher, W.W. (1991). Effects of an agonist of gonadotropin releasing hormone on ovarian follicle in cattle. J. Biol. Reprod. 45: 883-889.

Malvern, P.V. (1984). Pathophysiology of the puerperium. J. Dairy Sci. 72: 150-162.

Marion, G.G. and Gier, H.T. (1968). Factors affecting bovine ovarian activity after parturition. J. Anim. Sci. 27: 1621-1628.

McNatty, K.P. (1988). Reproductive endocrinology of postpartum cow. Proceeding. 5th Seminar Dairy Cattle Soc. N.Z. Vet. Assoc. P. 61.

Mialot, J.P.G.; Laumonnier, C. Ponsart, H.; Barassine, E.; Ponter, A.A. and Deletang, F. (1999). Postpartum subestrus in dairy cow: comparison of treatment with PGF<sub>2</sub> $\alpha$  or GnRH+ (PGF<sub>2</sub> $\alpha$  + GnRH). Theriogenology, 52: 911-917.

Michiel, G.; Bostedt, H.; Hoffman, B.; Failing, K. and Rattenberger, E. (1999). Effect of a 10 day postpartum administration of PGF<sub>2</sub> & analogue at different concentrations on the course of the puerperum and fertility in dairy cows. Tierarztl prax Ausg G. Grosstiere, 27: 16 – 24.

Montgomery, G.W. (1985). The effects of season on reproduction in beef cows. A review. Proc. J. Anim. Prod. 45: 47-48.

Moss, G.E; Parfect, J. R; Marvin, C.A; All rich, R.D and Dickman, M.A. (1985). Pituitary concentrations of gonadotropins and receptors for GnRH in suckled beef cows at ovarian intervals after calving J. Anim. Sci 60: 285 – 289.

Mukasa, M.E.; Tegene, A. and Ketema, H. (1991). Patterns of postpartum oestrus onset and associated plasma progesterone profiles in Bos indicus cows in Ethiopia. J. Anim. Reprod. Sci. 24: 73-84.

Olori, V.E. and Galesloot. P. (2000). Projection of partial lactations and calculation of 305-day yield in the republic of Ireland. Interpol Bull. No. 22: 149-154.

Olson, J.D.; Bal, L.; Mortimer, R.G.; Farin, P.W.; Aaney, W.S. and Huffrman, E.M. (1984). Affects of Bacteriology and endocrinology of cows with pyometra and retained fetal membranes. Amr. J.Vet. Res. 45: 2251-2255.

Osman A.H. and Eladin, F.M. (1971). Some dairy characteristics of Northern Sudan Zebu Cattle. 111 Reasons for Disposal of Dairy Cows. Trop. Agric. 48: 327-331.

Osman, A.H. (1984). Sudanese indigenous cattle breeds and the strategy for the conservation and improvement. FAO. Animal production and health paper. 44: 58-66.

Paul, D. (1995). Body Condition of the cow. Veoprs Holland, 23:12-13.

Peters, A.R. and Riley, G.M. (1982). Is the cow a seasonal breeder? Br. Vet. J. 138: 533-550.

Peters, A.R. (1984). Reproductivity of the cow in the postpartum period 1- Factors affecting the length of postpartum period. Bri. Vet. J. 140: 76-84.

Peters, A.R. and Lamming , G.E. (1984). Reproductive activity of the cow in the postpartum period. Endocrine patterns and induction of ovulation. Vet. J. 140: 259-280.

Peters, A.R. and Bosu, W.T.K (1988). Relationship of uterine infection and folliculogenesis in dairy cows during early puerperium. J. Theriogenology, 30: 1045 - 1055.

Pryce, J.E.; Veerkamp, R.F.; Thompson, R.; Hill, W.G.; and Simm, G. (1997). Genetic aspect of common health disorders and measures of fertility in Holstein Friesian dairy cattle. *Anim. Sci.* 65: 353-360.

Pryce, J.E.; Esslemont, R.J.; Thomposn. R.; Veerkamp, R.F.; Kossaibati, M.A. and Simm, G. (1998). Estimation of genetic parameters using health, fertility and production data from management recording system for dairy cattle. *Anim. Sci* 66: 577-584.

Pryce J.E; Coffey M.P and Smith, G. (2001). The relationship between body condition Score and reproductive performance. *J. Dairy Sci* 84: 1508 – 1515.

Randel, R.D. (1981). Effect of once daily suckling on postpartum interval and cow-calf performance of first calving in Brahman Hereford heifers. *Anim. Aci.* 53: 755-761.

Rhind, S.M. and McNeilly, A.S. (1986). Follicle population, ovulation rates and plasma profile of LH, FSH and prolactin in Scottish black face ewe in high and low level BCS. *Anim. Reprod. Sci.* 10: 105-115.

Rhodes, R.C.; Randel, R.D. and Long, C.R. (1982). Corpus luteum function in the bovine: in vivo and in vitro evidence for both a seasonal and breed type effect. *J. Anim. Sci.* 55: 159-168.

Richards, M.W; Wettelman R. P. And Schoenerman (1982). Nutritional oestrous in beef cows, body weight changes, B. C. L H in serum and ovarian activity. *J. Anim. Sci.* 67: 1520 – 1989.

Ruiz, F.; Magaly, G; Gonzalez, A.; and Fernandez, A.J. (1992) Reproductive performance of Zebu-Boss- Taurus, cross bred cows under an intensive management system. *Western Plairis. Anim. Breed. Abst.* 60: 288-289.

Savio, J.D; Boland, M.P.; Hynes, N and Roche, I.F (1990) Resumption of follicular activity in the early postpartum period of dairy cows. *J. of reprod. And fertility* 88: 569.

Schallenberger, E; Walters, D.L. and Oschman, R.S. (1984). Endocrine mechanisms contributing to postpartum period anoestrus in dairy cattle. *Proceding of the inter. Cong of Amin. Reprod. and A.I.* 10:206-220.

Schams, D.; Hofer, F.; Hofmann, B.; Ender, M.L. and Karg, H. (1973). Effects of synthetic L.H. R.H. treatment on bovine ovarian

function during oestrous cycle and postpartum period. *Acta. Endocrinol suppl.* 177: 296-311.

Schofield, S.A.; Kitwood, S.E and Philips, C.J. (1999). The effects of a postpartum injection of PGF2 $\alpha$  on return to oestrus and pregnancy rates in dairy cows, *Brit. Vet. J.* 157: 172 – 177.

Sejresan, K. and Sorensen, A. (1982). Nutritional physiology and feeding of the cow around parturition. *J. Anim. Sci.* 60: 325-334.

Short, R.E.; Bellow, E.L.; Staihgmler, R.B.; Berardinelli, J.G.; Custer, E.E. (1990). Physiological mechanism controlling anoestrus and fertility in postpartum beef cattle. *J. Anim. Sci.* 55: 699-816.

Smith, M.F.; Bueewl, W.C.; Ship, L.D.; Sprott, L.R.; Songster, W.N. and Wiltbank, J.N. (1987). Hormone treatments and calf removal in postpartum beef cows. *Anim. Sci.* 15: 12-22.

Snijders, S.E.M.; Dillon, P.G.; Farrell, K.J.O.; Diskin, M.; Wylie, A.R.G.; Callaghan, D. O.; Rath,M. and Boland, M.P. (2001). Genetic merit for milk production and reproductive success in dairy cows. *Anim. Reprod. Sci.* 65: 17-31.



Stagger, K.; Diskin, M.G.; Streeman, J.M.; Roche, J.F. and Piskin, M.G. (1998). Effect of calf isolation on follicular waves dynamic gonadotropin and metabolic hormone changes and interval to first ovulation in beef cows fed-either of two levels of energy postpartum. *J. Biol. Reprod.* 59: 777-783.

Staples, C.R.; Thather, W.W. and Clark, J.H. (1990). Relationship between ovarian activity and energy status during the early postpartum period of high producing dairy cows. *J. Dairy Sci.* 73: 938-947.

Stevenson, J. S.; Phatak, A. P; Rettmer, I. and Stewart, R.E (1993). Postinsemination administration of receptal : follicular dynamics duration of cycle , hormonal responses and pregnancy rates . *J. Dairy Sci.* 76: 25 – 36.

Strandberg, E. and Oltenacu, P.A. (1989). Economic consequences of different calving intervals. *Acta. Agric. Scand.* 39: 407-420.

Symington, R.B. and Hale, (1967). Methods to stimulate sexual activity in lactating Zebu cows. *Rhod. Zambia and Malawi. Agric. Res.* 5: 13-21.

Thatcher, W.W. (1974). Effects of season, climate and temperature on reproduction and Lactation. *J. Dairy. Sci.* 57: 360-368.

Thatcher . C.D. (1986). Effects of nutrition and management of the dry and fresh cow on fertility. *Bovine Pract.* 21: 172 - 179.

Thatcher, W.W; Driancourt, D. A. and Terquit, M. (1991) Dynamics of ovarion follicular development in cattle following hystrectomy and during early pregnancy. *Dom. Anim. Endocrinol*, 8: 228 – 224.

Thatcher, W.W.; Drost, J.D.; Savio, K.L.; Macmillan, R.L.; Delasota, M.S. and Morries, G.R. (1993). New clinical uses of GnRH and its analogues in cattle. *Anim. Reprod. Sci.* 33: 27-49.

Twagiramungu, H; Guilbault, L.A. and Dufour, J.J. (1995). Synchronization of ovarian follicular waves with a GnRH against to increase precrsion of oestrus in cattle. *J. Anim. Sci* 73: 3141-3151.

Vaccaro, L.P. (1974). Dairy cattle breeding in tropical South America. *World Animal Review.* 12: 8-13.

Veerkamp, R.F. (1998). Selection for economic efficiency of dairy feed intake. *J. Dairy Sci.* 81: 1109-1119.

Vicker, S.D.; Guire, W.J.; Wright, J.M.; Beeman, K.B. and Kircofe, G.H. (1989). Cow-calf association delays postpartum ovulation in mastectomized cows. *Theriogenology*, 32: 467- 471.

Villa-Godoy, A.; Hughes, T.L.; Emery, R.S.; Chapin, L.T. and Fogwell, R.L. (1988). Association between energy balance and luteal function in lactating dairy cows. *J. Dairy Sci.* 71: 1063-1072.

Webb, R.; Lamming, G.E.; Haynes, N.B.; Hafs, H.O. and Manns, J.G. (1996). Response of cyclic and postpartum suckled cows to injections of synthetic LH-RH . *J. Reprod. Fertil.* 50: 203-216.

Wildman, E.E.; Jones, G.M.; Wagner, P.E.; Boman, R.L.; Trout, H.F. and Ilesch, T.N. (1982) A dairy cow body condition scoring and its relationship to selected production characteristics. *J. Dairy Sci.* 65: 495-499.

William, G.L; Kotwiga, J.; slanger, W.D.; Olson, D.K.; Tilton, J.E. and Johnson, K.J. (1982). Effects of suckling on pituitary responsiveness to gonadotropin - releasing hormones throughout the early postpartum of beef cows. *J. Anim. Sci.* 54: 594 – 603.

Williams , G.L , 1990 sensory and behavioural control of gonadotropin secretion during suckling mediated anovulation in cows. J. Reprod . Fertil. (suppl) 49: 463- 475.

William, G.L. and Griffith, M.K. (1995). Sensory and behavioral control of gonadotropin secretion during suckling-mediated an ovulation in cows. J. Reprod. Fertil. (Supplement) 49: 463-475.

Wise, M.E.; Rodringuez, R.E.; Armstrong, D.V.; Huber, J.T.; Wiersma, F. and Hunter, R.E. (1988). Fertility and hormonal responses to temporary relief of heat stress in lactating dairy cow. Theriogenology, 29: 1027-1035.

Yavas, Y. and Walton, V. (2000). Postpartum cyclicity in suckled beef cows. Theriogenology, 54: 25-55.

Youngquist, M. A. (1988). Anoestrous and infertility in the cow In: *Fertility and infertility in Vet. Practice*. (4th ed.) Laing, J.A. (ed) Bailliere Tindall pp 91-112.

Zainelabdin, H.A (2001) *Some aspects of reproductive performance of Holstein Friesian cows in the Sudan* - M. Sc. Thesis, U of K.

Zeddies, J. (1982). Special economic aspects of fertility related to central European farming conditions. Current topics in Vet. Medicine and Anim. Sci. 20: 425-441.

Zemjanis, R. (1980). Repeat breeding in cattle. In: Current Therapy in Theriogenology. (1st ed.) P. 205-212.

## **Appendix .1.**

**The average of rain fall /per/ mm in Nyala station from 1999-2005**

**(Nyala Metrological office)**

<b>The year</b>	1999	2000	2001	2002	2003	2004	2005
<b>Nyala Station</b>	480.6	397.4	286.2	311.1	552.4	268.00	494.00

**Appendix .2.**  
**Population of livestock in South Darfur State.**

### **Appendix.3.**



## **Appendix .4.**

## **Appendix .5.**

### **Different parameters of PPI of local dairy cows native to South Darfur State**

<b>No</b>	<b>Breed</b>	<b>BCS</b>	<b>Parity</b>	<b>PPI</b>
1	Kenana	3	2	688
2	Kenana	3	2	485
3	Kenana	3	5	155
4	Kenana	3	2	313
5	Kenana	3	3	123
6	Kenana	3	3	92
7	Kenana	3	5	186
8	Kenana	3	2	273
9	Kenana	3	3	189
10	Kenana	3	3	180
11	Kenana	2.5	2	388
12	Kenana	3	2	268
13	Kenana	2.5	3	212
14	Kenana	2.5	3	152
15	Kenana	2.5	3	138
16	Kenana	3	3	138
17	Kenana	3	3	455
18	Kenana	3	5	360

19	Kenana	2.5	3	276
20	Kenana	2.5	3	203
21	Kenana	2.5	4	216
22	Fellata	2.5	5	203
23	Fellata	3	3	216
24	Fellata	2.5	3	221
25	Fellata	2.5	5	121
26	Fellata	2.5	3	124
27	Fellata	2.5	4	205
28	Fellata	2.5	3	345
29	Fellata	4	4	225
30	Fellata	3	3	310
31	Fellata	3	4	275
32	Fellata	3	3	210
33	Fellata	3	2	478
34	Fellata	3	3	274
35	Fellata	3	2	166
36	Fellata	3	5	138
37	Fellata	2.5	3	271
38	Fellata	3	3	203
39	Cross-bred	4	5	132
40	Cross-bred	3	2	184
41	Cross-bred	3	3	97
42	Cross-bred	3	2	229
43	Cross-bred	3	3	260
44	Cross-bred	3	3	74
45	Cross-bred	4	3	151
46	Cross-bred	3	3	140
47	Cross-bred	3	3	131
48	Cross-bred	3	4	71
49	Cross-bred	4	3	202
50	Cross-bred	3	1	173
51	Cross-bred	3	1	204
52	Cross-bred	3	2	152
53	Cross-bred	4	2	56
54	Cross-bred	4	1	44
55	Cross-bred	4	2	57
56	Cross-bred	4	2	70
57	Cross-bred	4	2	62

58	Cross-bred	4	1	44
59	Cross-bred	4	3	30

## Appendix .6.

### PPI of local breeds of dairy cows native to South Darfur State treated with GnRH

No	treatment	Kenana	Fellata	Cross-bred cows
1	treatment	76	58	86
2	treatment	60	77	84
3	treatment	90	63	63
4	treatment	31	64	77
5	treatment	79	55	71
6	treatment	30	80	83
7	treatment	58	57	64
8	treatment	57	69	68
9	treatment	111	55	60
10	treatment	126	85	81
11	treatment	107	62	79
12	Control	161	216	132
13	Control	377	221	184
14	Control	688	121	97
15	Control	485	124	229
16	Control	155	205	260
17	Control	313	345	74
18	Control	123	225	151
19	Control	92	510	140
20	Control	186	275	131
21	Control	273	210	71
22	Control	189	478	202

## Appendix .7.

**The length of PPI of local dairy cows native to South Darfur State during the three different seasons**

<b>No</b>	<b>Breed</b>	<b>Winter</b>	<b>Summer</b>	<b>Autumn</b>
1	Kenana	92	212	168
2	Kenana	161	152	277
3	Kenana	377	138	421
4	Kenana	688	476	155
5	Kenana	485	388	123
6	Kenana	155	76	189
7	Kenana	313	60	267
8	Kenana	123		180
9	Kenana	186		313
10	Kenana	276		
11	Kenana	189		
12	Kenana	180		
13	Kenana	386		
14	Fellata	203	225	138
15	Fellata	261	274	271
16	Fellata	121	166	203
17	Fellata	124	138	186
18	Fellata	205		273
19	Fellata	221		286
20	Fellata	345		
21	Fellata	478		
22	Fellata	210		
23	Fellata	510		
24	Fellata	275		
25	Cross-bred	151	30	132
26	Cross-bred	140	45	184
27	Cross-bred	131	33	97
28	Cross-bred	71	61	229
29	Cross-bred	202	45	
30	Cross-bred	173	42	
31	Cross-bred	204	67	

## Appendix .8.

**The effect of the season on the PPI of Kenana, Fellata and cross-bred dairy cows.**

<b>Breed</b>	<b>Summer</b>	<b>Winter</b>	<b>Autumn</b>
Kenana	214.57±60.00	276.39±45.9	233.56±31.8
Fellata	200.75±30.4	268.46±38.7	233.18±22.9
Cross-bred	46.00±6.5	153.14±17.4	160.00±28.9

## Appendix .9.

**Kenana breed dairy cows:-**

<b>The name of the cow</b>	<b>Parity</b>	<b>B.C.S</b>	<b>Day of part.</b>	<b>PPI</b>	<b>Day of conception</b>	<b>Repeat breeding</b>
WIZEAN	3	3	20.12.2001	161 day	30.5.2002	
ARDA	4	3.5	26.12.2001	377 day	7.2.2003	
DOWMA	2	4	20.12.2001	688 day	10.11.2002	
SAFRA	2	2.5	1.1.2002	485 day	1.5.2003	
BASHOM	5	2.5	5.1.2002	155 day	9.6.2002	Rep. Breeding 1.7.2002
ALTANMIN	2	3	6.1.2002	313 day	15.11.2002	
ARIAL	3	3	2.1.2002	123 day	5.7.2002	
BASKAWIT	3	2.5	16.11.2001	92 day	16.2.2002	
HORA	5	3	5.1.2002	186 day	10.7.2002	
HAMAN	2	2.5	5.1.2002	273 day	5.10.2002	
OMATIMAN	3	2.5	23.1.2002	189 day	31.7.2002	

OMATIMAN	3	3	5.2.2002	180 day	4.8.2002	
GARA	2	2.5	11.3.2002	388 day	3.2.2002	
BIT DOWMA	2	2.5	6.2.2002	368 day	1.2.2003	
SAFARA <sub>2</sub>	3	2.5	1.3.2002	212 day	3.8.2002	
KINANA	3	3	30.3.2002	152 day	1.9.2002	
DANG KOWAL	3	3	28.3.2002	138 day	16.8.2002	
HAMAM	5	3	4.4.2002	138	20.8.2003	
BKALIL	3	2.5	2.10.2003	455	1.4.2004	
KALL	3	2.5	20.10.2003	467	1.2.2004	
SAIRA <sub>2</sub>						
KENANA	4	2.5	10.1.2003	47.6	30.4.2004	

## Appendix .10.

### Fellata breed dairy cows

The name of the cow	Parity	B.C.S	Day of part.	PPI	Day of conception	Repeat breeding
KADRA	5	2.5	7.11.2001	2.3 day	29.4.2002	12.6.2002
BIKADR A	3	3	13.11.2001	216 day	2.8.2002	
OMRAS	3	2.5	6.12.2001	221 day	15.7.2002	
GALHA	5	2.5	5.11.2001	121 day	6.3.2002	
FERY	3	2.5	12.11.2001	124 day	16.3.2002	
BIT WIZIEAN	4	2.5	13.11.2003	205 day	5.6.2002	
ARNAB	3	2.5	15.12.2001	345 day	27.11.2002	
AL NIEA	4	4	20.3.2002	225 day	1.11.2003	
FEREY	3	3	8.11.2002	510 day	1.4.2004	
GILAHA	4	3	1.12.2002	275 day	3.10.2003	
ALTAHER	3	3	2.11.200	210 day	31.5.2003	

1			2			
ALTAHER 2	2	3	28.2.202	478 day	25.5.2003	
SAFRA	3	3	1.3.2003	274 day	30.11.2003	
OMGHIRA IEN	2	2.5				
OMGHIRAIEN	2	3	2.3.2003	166 day	15.8.2003	
HAMAM	5	3	4.4.2002	138 day	20.8.2003	
SIGAIRA	3	2.5	15.7.200 3	271 day	13.4.2004	
HAWA	3	3	20.7.200 3	203 day	10.2.2004	

## Appendix .11.

### Cross-bred:-

The name of the cow	Parity	B.C.S	Day of part.	PPI	Day of conception	Repeat breeding
ZARFGA	5	3.5-4	15.10.2001	132 day	26.2.2002	
GIRAYBA	2	3	13.12.2001	184 day	15.6.2002	
HAMAMA	3	3	20.12.2001	97 day	27.3.2002	
FORSA	2	3	23.12.2001	229 day	9.8.2002	
HORA	3	3	1.1.2002	260 day	17.9.2002	
OMGASH	3	3	20.12.2001	74 day	4.3.2002	
ZIRAG	3	3.5-4	1.1.2002	151 day	5.6.2002	
OMSAIER	3	3	25.12.2001	140 day	5.5.2002	
KARASI	3	3	27.1.2002	131 day	10.5.2002	
OMSAMGOR	4	3	25.2.2002	71 day	5.5.2002	
KOSTL	3	3.5-4	25.2.2002	202 day	15.8.2002	
GAZAL	1	3	28.2.2002	173 day	20.8.2002	
KAMSA	1	3	24.2.2002	204 day	21.9.2002	
SORA	2	3	13.4.2002	152 day	17.9.2002	
SALIGA	2	3.5-4	10.3.2002	56 day	5.5.2002	
BOLOWZA	1	3.5-4	22.4.2002	44 day	5.6.2002	
GADIDA	2	3.5-4	15.3.2002	57 day	12.5.2002	2.6
HEMAIER	3	3.5-4	17.3.2002	70 day	26.5.2002	Rep
TAHNIA	2	3.5-4	17.3.2002	62 day	3.5.2002	Rep
ONWAN	1	3.5-4	28.3.2002	44 day	11.5.2002	Rep 23.5
TAWILA	2	3.5-4	12.4.2002	30 day	12.2.2002	Rep 6.6.2002
MARYODA	4	3.5-4	5.4.2002	45 day	20.5.2002	Rep 8.6
TIKO	1	3.5-4	18.4.2002	33 day	21.5.2002	Rep
ISKARTO	1	3.5-4	25.4.2002			
GARNOG	1	3.5-4	22.3.2002			



OMMASIER	3	3.5-4	2.4.2002			
BREAK	3	3.5-4	24.4.2002			
GARA	2	3	24.4.2002			
BIT HALOF	2	3	1.5.2002			
ZARAF	3	3				

## **Appendix .12.**

**Kenana dairy cows treated with (GnRH) :-**

<b>The name of the cow</b>	<b>Parity</b>	<b>B.C.S</b>	<b>Day of part.</b>	<b>PPI</b>	<b>Day of conception</b>	<b>Repeat breeding</b>
GARA	3	3	14.3.2002	76 day	20.6.2002	
RIDINA	5	3.5-4	23.3.2002	60 day	11.6.2002	
WIZEAN	3	3	29.9.2002	90 day	20.1.2003	
OMMARA D	4	3.5-4	30.9.2002	31 day	31.10.2003	
BITWIZE AN <sub>A</sub>	3	3	16.2.2003	80 day	30.5.2003	
BITWIZA N <sub>B</sub>	2	3	13.12.2002	58 day	9.2.2003	
SOMEAT	3	3	20.12.2002	57 day	15.2.2003	
HINAKE	4	3	7.3.2003	111 day	27.7.2003	
ASAL	2	3	29.1.2003	126 day	26.6.2003	Rep. Br.
MOZA	3	3	2.7.2002	107 day	12.11.2003	
BASHOM	4	3	4.4.2003	76 day	10.7.2003	
KINANIA	5	3	17.2.2003	89 day	6.7.2003	
ASAL <sub>2</sub>	3	3	15.6.2003	106 day	24.10.2003	

## **Appendix .13.**

**Fellata dairy cows treated with GnRH.**

<b>The name of the cow</b>	<b>Parity</b>	<b>B.C.S</b>	<b>Day of part.</b>	<b>PPI</b>	<b>Day of conception</b>	<b>Repeat breeding</b>
BIT WIZEAN	5	3	10.4.2003	58 day	7.6.2003	
WIZA	2	3.5-4	7.4.2003	77 day	23.6.2003	
KADRA	5	3.5-4	27.4.2003	63 day	30.6.2003	
GMRAS	4	3	6.5.2003	64 day	9.7.2003	
BIT KADRA	4	3	8.5.2003	55 day	2.7.2003	
SIGAIERA	2	3	7.8.2003	80 day	27.10.2003	
HAWAYA	3	3.5-4	14.8.2003	57 day	11.10.2003	
HAWAYA	2	3.5-4	26.8.2003	69 day	4.11.2003	
ARNAB	4	3.5-4	13.9.2002	55 day	17.11.2003	
GALHA	5	3	7.10.2003	85 day	13.12.2003	
FEREY	4	3	8.1.2003	62 day	10.3.2003	
	5					

**Appendix .14.**  
**Cross-bred dairy cows treated with GnRH.**

<b>The name of the cow</b>	<b>Parity</b>	<b>B.C.S</b>	<b>Day of part.</b>	<b>Day of injection</b>	<b>PPI</b>	<b>Day of conception</b>	<b>Repeat breeding</b>
GIRAYPA	3	3	21.7.2003	15.8.2003	86 day	16.10.2003	
ADILA	2	3	9.7.2003	1.8.2003	84 day	2.10.2003	
BIT ZARAF	2	3	8.10.2003	29.10.2003	63 day	10.12.2003	
GINANA	3	3	7.9.2003	29.9.2003	77 day	23.11.2003	
GAZAL	4	3	12.10.2003	2.9.2003	71 day	22.12.2003	
KOSTI	3	3	1.11.2003	22.11.2003	83 day	22.1.2004	
OM SANGOR	5	3	17.11.2003	8.12.2003	64 day	20.1.2004	
ZARAL	4	3	18.11.2003	9.12.2003	68 day	25.1.2004	
SOMEAT	3	3	18.11.2003	9.12.2003	60 day	17.1.2004	

SALIGA	2	3	20.11.2003	11.12.2003	81 day	9.2.2004	
GARA	2	3	12.12.2003	2.1.2004	79 day	29.2.2004	
TAHNIA	2	3	17.3.2002		62 day	3.5.2002	Rep
BOLOZA	1	4	22.4.2002		44 day	5.6.2002	Rep